

Page numbers followed by *f* indicate figures and those followed by *t* indicate tables.

A

Acceleration detection, 19
 Acceleration performance, and predator attack outcomes, 79
 Acceleration reaction, 7
 Accommodation, 27–28
 Acoustic perception, 23–24, 25f. *See also* Hydrodynamic perception
 Age-0 bluegills. *See also* Juveniles; Larval bluegills
 body depth of, 77, 78f
 early growth phases of, 119
 length–mass relationship of, 119
 survivorship of, 120–121, 121f
 Age at maturity, 87
 Age determination, 119
 Alewives (*Alosa pseudoharengus*), 22
 Alkalinity, and growth, 128
Ambloplites rupestris, 70, 70f, 71
 Ambush predators, 79, 82
Ameiurus natalis, 70, 71
Ameiurus spp., as nest predators, 102, 105
 Amphibians, behavioral alterations due to predator presence, 73
 Androgens, 99
 Angling, impact of, 141–143, 142f, 145–146
 Antimycin A, 141
 Apparent size, defined, 43
 Apparent size model (ASM), 43–46, 44f, 45t, 46f, 49–50
 Aquatic vegetation. *See* Vegetation
 ASM. *See* Apparent size model (ASM)
 Auditory threshold, 24, 25f

B

Background spacelight, 30
 Banded puffer (*Colomesus psittacus*), 21, 21f
 Behavioral thermoregulation model (BTM), 42–43
 Biological control of vegetation, 136, 147
 Black crappie (*Pomoxis nigromaculatus*), 70, 70f, 71
 Bluegill-pumpkinseed hybrids, reproductive behavior of, 98, 99f, 100f
 Body depth
 of age-0 bluegills, 77, 78f
 vs. body length, 75, 76f
 and predation, 75, 77, 81
 Body form, and predator attack outcomes, 78–79
 Body length
 body depth and, 75, 76f
 body mass and, 119, 124, 125f
 vs. scale length, 119f
 visual acuity and, 24–25, 26f, 27f, 31–32, 32f
 Body mass, vs. body length, 119, 124, 125f
 Bound circulation, 3
 Bound vortex, 3

Bourgeois males. *See* Territorial males
 Braking, 15–17, 17f
 Breeding. *See* Reproduction
 Breeding age, 87
 Breeding season, 87–88, 88f
 BTM (behavioral thermoregulation model), 42–43
 Buccal pressure, during feeding, 36, 37f
 Bullheads (*Ameiurus* spp.), 70, 71, 102, 105
 Buoyancy, center of, 13

C

Cannibalism, 103, 114
 Captivity, artifacts of, 84, 133
 Catch limits, 142
 Caudal fin
 in braking, 17
 in nest sweeping, 91
 in propulsion, 11–12, 12f, 13f
 Center of buoyancy, 13
 Center of mass, 10, 15
 Channeling of vegetation, 137
Chloeon dipterum, 23, 23f
 Circulation, 3
Colomesus psittacus, 21, 21f
 Colonial nesting, 89, 89f, 104–105, 104f
 Coloration
 cuckolds, 108
 female bluegills, 91
 male bluegills, 90, 92, 108
 pumpkinseeds, 97
 Common carp (*Cyprinus carpio*), 132
 Community dynamics, foraging and, 56–58
 Competition, 59–73
 interspecific, and dietary shifting, 64–69, 64t, 65t, 66t, 68t, 72
 intraspecific, 69–70
 invertebrates and, 71
 and juvenile habitat/diet shifts, 59–64, 60t, 61t, 62–63f, 63t
 lack of, and disruption of ontogenetic diet shifts, 71–72
 of less desirable fish, 143–145, 145f
 piscivores and, 63–64, 63t, 70–71, 70f
 prey behavior alterations and, 72–73
 zebra mussel and, 72
 Cones, 24–25, 27, 27f, 30–31
 Conspecific recognition, 38, 102
 Contrast sensitivity, 31
 Courtship
 behavioral repertoires, 94, 94t
 female behavior, 91–92
 male behavior sequences, 95–98, 95–100f
 sequential behaviors in, 92, 93f
 terminology, 93t

- Crayfish, 136, 147
Ctenopharyngodon idella, 136, 147–148
Cuckolders
 breeding success of, 111–112
 life span of, 109
 relative numbers of, 109, 146
 reproductive strategy of, 108, 108f
 sperm characteristics of, 110–111, 111f
 and stunting, 141
 testes mass of, 112
Cuckoldry, 107–115
 fitness of male polymorphs, 107, 109–115, 110f, 111f, 113f
 reproductive strategies of male polymorphs, 107–109, 108f
Cupulae of lateral line, 19, 21
Cyprinus carpio, 132
D
Daily feeding patterns. *See* Diel foraging
Damsselfly larvae (*Enallagma cyathigerum*), 22–23, 23f
Daphne lumholtzi, 48, 49f
Daphne magna, 48, 49f
Daphnia. *See* Water fleas (*Daphnia*)
Daylength, and gonadosomatic index, 88, 88f
Delayed maturation, 87
Depth, body. *See* Body depth
Detached eddy, 3
Development, 117–119, 118f, 119f
Diel foraging, 50–53, 51f, 52t
Dietary constituents, 33–34, 50, 52t
 for juveniles, 59–60, 60t, 61t
 for larvae, 118
Dietary overlap
 with green sunfish, 67–69, 68t
 with pumpkinseeds, 64t, 65–67, 65t, 66t
 with redspotted or redear sunfish, 72
Dietary shifting
 and growth, 128
 interspecific competition and, 64–69, 64t, 65t, 66t, 68t, 72
 by juveniles, 59–64, 60t, 61t, 62–63f
Diet width, habitat complexity and, 38
Digestion rate, temperature and, 127–128
Displacement detectors, 19
Dorosoma cepedianum. *See* Gizzard shad (*Dorosoma cepedianum*)
Dorosoma petenense, 137, 147
Dorsal fin, 11–12, 11f, 12f, 13f, 17
Drag propulsion, 7, 8f
Dreissena polymorpha, 72
E
Eccritic temperature, 126
Eggs
 development of, 117–119, 118f
 Eggs (*cont.*)
 number per nest, 92
 predation on, 102–105, 104f, 106f
Enallagma cyathigerum, 22–23, 23f
Encounter rate, 79
Energy balance
 foraging behavior and, 40–41, 45, 48
 and growth, 126
Environmental factors, and growth, 128
Epiphytic invertebrates
 snags and, 55–56
 vegetation and, 53–55, 55t
Escape of prey, 48
Esocids, prey selection by, 79–80
Esox lucius. *See* Northern pike (*Esox lucius*)
Esox masquinongy, 22
Eurasian water milfoil (*Myriophyllum spicatum*), 133, 134
European catfish (*Silurus glanis*), 21
Excretory products, 58
F
Fanning of nest, 99–100
Feeding, water temperature and, 123, 126–127, 127f
Feeding adaptations, 34–38, 34–37f. *See also* Foraging
Feeding rate, 39
Female mimics. *See* Satellites
Females
 courtship/spawning behavior of, 91–92, 115
 male preference by, 115
 nest predation by, 103
Fertilizers, 147–148
Fin rays, and predator attack outcomes, 78–79, 80
Five-spot African cichlid (*Thysochromis ansorgii*), 21, 21f
Food availability
 diet width and, 38
 and growth, 128
Foraging, 33–58
 diel patterns, 50–53, 51f, 52t
 dietary preferences, 33–34, 50, 52t
 feeding adaptations, 34–38, 34–37f
 and lake community dynamics, 56–58
 littoral to limnetic shift of juveniles, 59–64, 62f, 63f
 models of. *See* Foraging models
 structural complexity and, 53–56, 55t, 57f
Foraging models, 38–50
 apparent size model, 43–46, 44f, 45t, 46f, 49–50
 behavioral thermoregulation model, 42–43
 diet width and, 38
 greatest stimulus model, 47–48, 49f
 handling time and, 38–39
 informed decision model, 47
 optimum foraging model, 40–42, 41f, 49–50
 prey characteristics and, 39–40
 reaction field volume model, 46–47
Frogs, behavioral alterations due to predator presence, 73
Fungal infections, and egg loss, 105, 106f

G

Gap, 94t
Gape
 of largemouth bass, 75, 76f
 limitation of prey size by, 75, 77, 81
Gill rakers, 34–35, 35f
Gizzard shad (*Dorosoma cepedianum*), 79–80, 137, 143–145, 145f, 147
"golden eye" stage, 118
Gonadal changes, 87, 88, 88f
Gonadosomatic index (GSI), 88f, 112, 140f
Grass carp (*Ctenopharyngodon idella*), 136, 147–148
Greatest stimulus model (GSM), 47–48, 49f
Green-sensitive cones, 31
Green sunfish (*Lepomis cyanellus*)
 competition by, 62, 65, 67–69, 68t
 habitat preferences of, 137–138
 prey detection by, 22
Growth, 119–129
 by age class, 122–123, 122f
 of earlier vs. later hatches, 120–121, 121f
 environmental factors and, 128
 interlake differences in, 123–124
 nutritional balance and, 124–126, 124t, 125f
 predation and, 128–129
 rapid/slow phases in age-0 bluegills, 119
 seasonal variations in, 126–127, 127f
 stocking levels and, 119–120
 water temperature and, 123, 126–128
GSI. *See* Gonadosomatic index (GSI)
GSM (greatest stimulus model), 47–48, 49f
Guppy (*Poecilia reticulata*), 21

H

Habitat–diet relationships, 38
Habitats, profitability of, 40, 69
Habitat shifts, by juveniles, 59–64, 62f, 63f
Habitat structural complexity
 and diet width, 38
 foraging and, 53–56, 55t, 57f
 management of, 137–139
 predation evasion and, 82–84, 85f
Habituation, 101–102
Hair cells, 19
Handling time
 bluegills vs. pumpkinseeds, 35, 36f
 defined, 35
 foraging efficiency and, 38–39
 piscivores, 79, 80, 81
Hatching, 117
Hatch survivorship/growth, spawning time and, 120–121, 121f
Hearing, 23–24, 25f
Herbicides, 147
Home ranges, 42
Homing, 42

Horn's index, 72
Hovering, habitat structural complexity and, 56, 57f
Hybridization, 106–107, 107t
Hybrids
 parental care by, 115
 reproductive behavior of, 98, 99f, 100f
Hydra canadensis, 103
Hydrodynamic perception, 19–23, 20f, 21f, 23f. *See also*
 Acoustic perception
Hydrodynamic principles, 3, 4f
Hydrodynamic trails of prey fish, 21, 21f

I

IDM (Informed decision model), 47
Inertial suction, in feeding, 35–36, 37f
Informed decision model (IDM), 47
Inner ear
 in acceleration detection, 19
 hair cells of, 19
 otoliths of, 19, 22
 semicircular canals of, 19
 sensory functions of, 19, 23
 swim bladder bridges to, 24
Insects/insect larvae
 behavioral alterations due to predator presence, 72–73
 detection of, 22–23, 23f
 in diet, 34–35
Intercone spacing, 24–25, 27, 27f
Interspecific competition
 dietary shifting and, 64–69, 64t, 65t, 66t, 68t, 72
 piscivores and, 63–64, 63t, 70–71, 70f
Intraspecific competition, 69–70
Invertebrates
 competition by, 71
 snags and, 55–56
 vegetation and, 53–55, 55t

J

Juveniles. *See also* Age-0 bluegills; Larval bluegills
 diet size selectivity of, 59
 feeding parameters of, 41–42
 habitat/diet shifts by, 59–64, 60t, 61t, 62–63f, 63t
 as nest predators, 102
 visual acuity of, 24–25, 26f, 31–32, 32f

K

Kelvin's theorem, 3

L

Labriform swimming, 4
Labyrinth. *See* Inner ear
Lake community dynamics, foraging and, 56–58
Largemouth bass (*Micropterus salmoides*)
 competition by, 63–64, 63t, 70, 70f, 71
 gape size of, 75, 76f, 77–78, 78f
 gizzard shad competition, 144, 145f

- Largemouth bass (*Micropterus salmoides*) (*cont.*)
 predation by, in structurally complex habitats, 82
 predation by, and bluegill quality, 131–132, 135
 prey detection by, 22
- Larval bluegills. *See also* Age-0 bluegills; Juveniles
 age determination, 119
 development of, 117–119, 118f
 identification of, 117
 predation on, 102–105, 104f, 106f
 survival of, adult population levels and, 69
 zebra mussel competition and, 72
 zooplankton populations and, 69
- Lateral display, 94t, 97
- Lateral line
 acoustic perception by, 19–20
 cupulae of, 19, 21
 hair cells of, 19
 in hydrodynamic detection, 19
 neuromasts of, 19, 21
 in spatial difference detection, 19, 20
 velocity detection by, 19, 21
- Lateral line canal, in velocity/acceleration detection, 19
- Leap, 94, 94t
- Length. *See* Body length
- Lens diameter, 25
- Lepisosteus oculatus*, 82
- Lepomis cyanellus*. *See* Green sunfish (*Lepomis cyanellus*)
- Lepomis gibbosus*. *See* Pumpkinseed (*Lepomis gibbosus*)
- Lepomis megalotis*, 137, 138
- Lepomis microlophus*, 72, 138
- Lepomis miniatus*, 72
- Levin's index, 65t, 66
- Lift propulsion, 7, 8f
- Light intensity
 foraging behavior and, 47
 reaction distance and, 28
 vision and, 28, 31
- Light penetration, vegetation and, 133
- Littoral to limnetic foraging shift of juveniles, 59–64, 62f, 63f
- Locomotion. *See* Propulsion
- Longear sunfish (*Lepomis megalotis*), 137, 138
- M
- Macroinvertebrates, habitat structural complexity and, 53–56, 55t
- Macrophytes. *See* Vegetation
- Male contingent reproductive strategies. *See* Cuckoldry
- Male polymorphs
 fitness of, 107, 109–115, 110f, 111f, 113f
 reproductive strategies of, 107–109, 108f
 and stunting, 139–141, 140f
- Male reproductive behavior repertoires
 bluegill-pumpkinseed hybrids, 98, 99f, 100f
 bluegills, 96–98, 97f, 98f
 pumpkinseed, 95–96, 95f, 96f
- Management, 131–148
 angling, 141–143, 142f
 bluegill quality, predation and, 131–132, 135
 competition of less desirable fish, 143–145, 145f
 habitat structure, 137–139
 ideal conditions, 145–148
 stunting, 139–141, 140f
 vegetation, 132–137, 135t
 water quality, 132
- Mass. *See* Body mass
- Maturity, age at, 87
- Mayfly larva (*Chloeon dipterum*), 23, 23f
- Median fins
 in braking, 17
 in movement, 8–13, 9–14f
 in nest sweeping, 91
- Micropterus salmoides*. *See* Largemouth bass (*Micropterus salmoides*)
- Milt, 110–111
- Morphological variation. *See* Polymorphism
- Mortality
 adults, 129–130, 129t, 146
 angling and, 143
 juveniles, 120–121, 121f
- Mouth, 34–36
- Movement, 3–17
 braking, 15–17, 17f
 hydrodynamic principles, 3, 4f
 median fins, 8–13, 9–14f
 pectoral fin kinematics, 4–7, 5f, 6f, 8, 9f
 propulsion, 7–12, 8f, 10f, 11–13f
 turning, 13–15, 16f
 wake dynamics, 3–7, 5f, 6f, 10f
- Mowing of channels, 137
- Muskellunge (*Esox masquinongy*), 22
- Myriophyllum spicatum*, 133, 134
- N
- Narrow diet hypothesis, 38
- Navigation, 42
- Negative stimuli, foraging behavior and, 48
- Nest colonies, 89, 89f, 104–105, 104f
- Nests
 abandonment of, 114
 building of, 90–91
 defense of, 98–99, 100–102, 102t, 104f, 113–114
 fanning of, 100–102
 site selection for, 89–90
 size of, 91
- Neuromasts, 19, 21
- Nitrogen excretion, 58
- Nitrogen supplementation, 148
- Nonselective predation, and growth, 129
- Northern pike (*Esox lucius*)
 attack tactics of, 79, 82
 competition by, 70, 70f, 71

Northern pike (*Esox lucius*) (*cont.*)

- prey selection by, 80
- size of bluegills preyed upon, 75

Nutrient enrichment, 147–148

Nutritional balance, and growth, 124–126, 124t, 125f

O

OFM. *See* Optimum foraging model (OFM)

Oncorhynchus mykiss, 15, 16

Opercular pressure, during feeding, 36, 37f

Opercular spread, 94t, 95, 96, 97

Opercular tab coloration in territorial males, 90

Optimal foraging rate, 40, 41f

Optimum foraging model (OFM), 40–42, 41f, 49–50

Otoliths, 19, 22, 119

P

Parental care, 98–102, 102t

Paternity, 111, 113–114, 113f

Pectoral fins

- in braking, 15–17, 17f
- kinematics of, 4–7, 5f, 6f, 8, 9f
- in propulsion, 7–8, 13f
- in turning, 14–15, 16f

Perca flavescens, 70, 70f

Pharyngeal pads, 34, 34f

Pharyngeal sound production, 94, 94t, 95–96

Phosphorus excretion, 58

Phosphorus supplementation, 148

Phytoplankton, 148

Piscivores. *See also names of specific species*

- active hunters vs. ambushers, 79
- in interspecific competition, 63–64, 63t, 70–71, 70f

Plants. *See also* Vegetation

- in diet, 33, 34, 50
- exotic, 133, 134, 136
- macroinvertebrates and, 53–55

Poecilia reticulata, 21

Polymorphism

- habitat structure and, 56, 62–63
- of male bluegills, 107–109, 108f. *See also* Cuckoldry
- predation and, 81

Pomoxis nigromaculatus, 70, 70f, 71

Populations, size structures of, 139–141

Predation, 75–85

- artifacts of captivity in studies of, 84
- and bluegill quality, 131–132, 135
- on eggs and larvae, 102–105, 104f, 106f
- evasive behaviors in avoiding, 79–84
- and growth, 128–129
- habitat complexity and, 138–139
- handling time and, 79, 80, 81
- polymorphism and, 81
- predator gape vs. body depth and, 75–78, 76f, 77f, 78f
- soft vs. spiny fin rays and, 78–79
- vegetation density and, 81–85, 85f

Predators. *See also* Piscivores

- detection of, by prey species, 72–73
- of nests, 102–103
- prey detection by, 20–23

Prey

- apparent size of, 43, 44f, 46f
- behavioral alterations due to predator presence, 72–73, 82–83
- characteristics influencing outcome of predator attacks, 78
- characteristics of, and foraging, 39–40, 47–50
- distinctive hydrodynamic trails of, 21, 21f
- escape of, 48
- evasive behaviors of, 49–50, 79–84
- habitat structural complexity and, 53–56
- movement of, 47
- predator detection of, 20–23
- transparency of, and reaction distance, 39–40

Prey encounter rate, 39

Propulsion, 7–8, 10f, 13f

- median fins in, 8–12, 11f, 12f
- pectoral fins in, 7–8, 8f

Protein balance, growth and, 124t, 125–126, 125f

Pumpkinseed (*Lepomis gibbosus*)

- breeding season of, 88
- competition by, 64–68, 64t, 65t, 66t
- diel foraging of, 50, 51f
- epiphytic invertebrate foraging by, 54
- gill rakers of, 35, 35f
- growth vs. prey size, 40–41
- habitat preferences of, 137–138
- handling times for, 35, 36f
- hydrodynamic trails of, 21, 21f
- male reproductive behavior, 95–96, 95f, 96f
- nest building, 91
- nesting of, 89–90, 105
- pharyngeal pads of, 34, 34f
- sequential courtship behaviors, 92, 93f

Q

Quality management. *See* Management

Quitting time, 42, 48–49

R

Rainbow trout (*Oncorhynchus mykiss*), 15, 16

Ram feeding, 36–37

Reaction distance (RD)

- defined, 25–26
- light intensity and, 28, 31
- prey transparency and, 39
- size-related probability of prey capture and, 44–45, 45t

Reaction field volume model (RFVM), 46–47

Redear sunfish (*Lepomis microlophus*), 72, 138

Red-sensitive cones, 31

Redspotted sunfish (*Lepomis miniatus*), 72

Relative vulnerability, 77

- Repertoire, defined, 94
Reproduction, 87–115. *See also* Courtship; Spawning
 breeding season, 87–88, 88f
 hybridization, 98, 99f, 100f, 106–107, 107t, 115
 male contingent strategies. *See* Cuckoldry
 maturity, 87
 nest building, 90–91
 nest site selection, 89–90, 89f
 parental care, 98–102, 102t, 104f, 113–114
 predation on eggs and larvae, 102–105, 104f, 106f
Resident males. *See* Territorial males
Retina, 24. *See also* Cones
Reverse von Kármán vortex street, 3, 4f
RFVM (reaction field volume model), 46–47
Ricco's law, 31
Rock bass (*Ambloplites rupestris*), 70, 70f, 71
S
Salamanders, behavioral alterations due to predator
 presence, 73
Saline environments, adaptation to, 35
Sander vitreum, 80
Satellites, 108, 108f. *See also* Cuckolders
Scales, in age determination, 119, 119f
Schoener's index, 66, 66t, 67, 68
Schooling, 82–83
Seasonal variations, in feeding and growth, 126–127, 127f
Sedimentation, vegetation and, 134
Semicircular canals, as angular acceleration detectors, 19
Sensory perception. *See* Acoustic perception; Hydrodynamic
 perception; Vision
Sexual dimorphism in males. *See* Male polymorphs
Shape, and propulsion, 8
Sight. *See* Vision
Silurus glanis, 21
Size, relative, in predation, 81
Size limits, 142, 143
Size-selective predation, and growth, 128
Snags, epiphytic invertebrates and, 55–56
Snails (*Viviparous georgianus*), 102, 105
Sneakers, 108. *See also* Cuckolders
Solitary nesting, 90, 105
Sound
 detection of. *See* Hearing
 high-intensity, and auditory threshold, 24, 25f
 low-frequency, lateral line detection of, 19–20
Spatial differences, lateral line detection of, 19, 20
Spatial summation, visual, 31
Spawn formation, 94t
Spawning. *See also* Courtship
 described, 91–92
 seasonal onset of, 87–88
 time of, and hatch survivorship/growth, 120–121, 121f
Sperm characteristics, 110–111, 111f
Sperm competition, 110–112, 110f
Sphere, submerged vibrating, dipole field of, 20, 20f
Spotted gar (*Lepisosteus oculatus*), 82
Starting vortex, 3
Stocking levels, and growth, 119–120
Structural complexity. *See* Habitat structural complexity
Stunting
 angling and, 143, 146
 causes of, 139–141, 140f
 heavy vegetation growth and, 133
Suction, in feeding, 35–36, 37f
Survivorship, of hatch, 120–121, 121f. *See also* Mortality
Swim bladder, in acoustic perception, 24
Swimming speed, 4, 78
Switch size, 61–62, 62f
T
Tail, in movement, 8–9, 10f
Temperature
 eccritic, 126
 and foraging area selection, 42–43
 and gonadosomatic index, 88, 88f
 and growth, 123, 126–128
Territorial maintenance, 94t. *See also* Nest, defense of
Territorial males
 angling and, 142–143
 cannibalism and nest abandonment among, 113–114
 early maturation of, 140, 140f, 143
 life span of, 108–109
 opercular tab coloration of, 90
 reproductive strategy of, 107
 sperm characteristics of, 110–111, 111f
 testes mass of, 112
Testes mass, 88t, 112. *See also* Gonadosomatic index (GSI)
Thermal stratification, 62
Threadfin shad (*Dorosoma petenense*), 137, 147
Thysochromis ansorgii, 21, 21f
Total length. *See* Body length
Turbidity
 control of, 147
 foraging behavior and, 47
 increased, effects of, 132
 vegetation and, 134
 vision and, 28, 31
Turning, 13–15, 16f
V
Vegetation. *See also* Plants
 and bluegill quality, 132–134
 density of, and predation, 81–85, 85f
 effects of, 134–136, 135f
 epiphytic invertebrates and, 53–55, 55t
 in habitat complexity, 138
 management, 132–137, 135t
 removal of, 136–138, 147, 148
Veiling radiance, 30
Velocity detection, by lateral line, 19, 21

Vision, 24–32

- apparent size model and, 44–45
- body size and, 24–26, 26f, 27f, 31–32, 32f
- vs. hydrodynamic perception in prey detection, 22
- light intensity/turbidity and, 28, 31
- position of prey and, 27, 29f
- spatial summation and, 31
- visual pigments and, 30–31

Visual acuity, 24–25, 26f, 27f

Visual angle, 24–26, 27f

Visual appearance, in habituation, 102

Visual pigments, 30–31

Visual stimuli, foraging behavior and, 47–48

Visual volume, 26, 28f

Viviparous georgianus, 102, 105

von Kármán vortex street, 3, 4f

Vortex rings, of wake, 4–7, 5f, 6f

Vortices

- starting vs. bound, 3
- submerged vibrating sphere model of, 20, 20f

Vorticity, total, 3

Vulnerability, relative, 77

W

Wake attenuation, 20–21

Wake dynamics, 3–7, 5f, 6f, 10f

Wakes, predator detection of, 21–23

Walleye (*Sander vitreum*), 80

Water fleas (*Daphnia*)

- pigmentation of, and feeding, 39–40
- spatial distribution of populations, 64

Water quality, 132, 147

Water temperature. *See* Temperature

Weberian ossicles, 24

Weber's law, 31

Wels (*Silurus glanis*), 21

Wide diet hypothesis, 38

Y

Yellow bullhead (*Ameiurus natalis*), 70, 71

Yellow perch (*Perca flavescens*), 70, 70f

Yolk sac larvae, 117

Z

Zebra mussel (*Dreissena polymorpha*), 72

Zooplankton feeding, switch to, 61, 62f, 64

Zooplankton populations

- bluegill feeding and, 57–58
- and bluegill quality, 132
- larval feeding and, 69
- phytoplankton and, 148